

#### **SAVE A LIFE**

## STOP THE BLEED® Course American College of Surgeons

Copyright © 2019 American College of Surgeons

Version 2



STOPTHEBLEED.ORG













The American
College of Surgeons
Committee on
Trauma

The American
College of
Emergency
Physicians

The National
Association of
Emergency Medical
Technicians


The Committee on Tactical Combat Casualty Care

# Some of the images shown during this presentation may be disturbing to some people.

## Why Do I Need This Training?

The #1 cause of preventable death after injury is bleeding.

## Where Can I Use This Training?



#### Goals

## 1. Identify

Recognize life-threatening bleeding

## 2. Stop the Bleed

Take steps to STOP THE BLEEDING

- **√** Pressure
- ✓ Packing
- **√** Tourniquets

## **Personal Safety**

## **YOUR** safety is **YOUR** first priority

- If you are injured, you cannot help others
- Help others only when it's safe to do so
- If the situation changes or becomes unsafe:
  - √ Stop
  - √ Move to safety
  - √ If you can, take the victim with you

## **Personal Safety**

#### **YOUR** safety is **YOUR** first priority

- Wear gloves if you can
- If you get blood on you, be sure to clean any part of your body that the blood has touched
- Tell a health care provider that you got blood on you, and follow his or her direction

A Alert 911

**B** Bleeding

**C** Compress

- A Alert 911
- **Bleeding**
- Compress



## A Alert 911

- Call 911
- Know your location
- Follow instructions provided by 911 operator

A Alert 911

**B** Bleeding

Compress



## **B** Bleeding


- Find source of bleeding
- Look for:
  - √ Continuous bleeding
  - √ Large-volume bleeding
  - √ Pooling of blood


## **B** Bleeding


- There may be multiple places the victim is bleeding
- Clothing may also hide life-threatening bleeding

## **B** Bleeding

- Arms and legs
- Neck, armpits, and groin
- Body







A Alert 911

**Bleeding** 

C Compress - Pressure




## C Compress - Pressure

- Apply direct pressure to wound
- Focus on the location of the bleeding
- Use just enough gauze or cloth to cover injury
- If pressure stops the bleeding, keep pressure on wound until help arrives



## C Compress - Packing


- For large wounds, superficial pressure is not effective
- If bleeding is from a deep wound, pack gauze tightly into the wound until it stops the bleeding; hold pressure until help arrives







## C Compress - Packing

- Arms and legs
- Neck, armpits, and groin
- Body







## C Compress - Tourniquet

- Apply 2 to 3 inches above wound
- Do not place over the elbow or knee
- Tighten tourniquet until bleeding stops
- Do NOT remove the tourniquet

## **C** Compress - Tourniquet

- Can apply to others or on yourself
- Can be applied over clothes
- Tourniquets HURT
- A second tourniquet may be required to stop the bleeding













## **CoTCCC Recommended STB Tourniquets**

#### **Recommended Non-Pneumatic Limb Tourniquets**

- Combat Application Tourniquet Gen 6 (CAT-6)
- Combat Application Tourniquet Gen 7 (CAT-7)
- Ratcheting Medical Tourniquet (RMT) Tactical
- SAM Extremity Tourniquet (SAM-XT)
- SOF Tactical Tourniquet—Wide (SOFTT-Wide)
- Tactical Mechanical Tourniquet (TMT)
- TX2 Tourniquet (TX2)
- TX3 Tourniquet (TX3)

## **CoTCCC Recommended STB Tourniquets**

#### **Recommended Pneumatic Limb Tourniquets**

- Delphi EMT (EMT)
- Tactical Pneumatic Tourniquet 2" (TPT2)

## Bleeding control in children

- In all but the extremely young child, the same tourniquet used for adults can be used in children.
- For the infant or very small child (tourniquet too big), direct pressure on the wound as described previously will work in virtually all cases.
- For large, deep wounds, wound packing can be performed in children just as in adults using the same technique as described previously.

#### **FAQs**

- Impaled objects?
- Improvised tourniquets?
- Loss of arm or leg?
- Pain?
- Other questions?

## **Summary**

- ✓ Personal safety
- A Alert 911
- **B** Find bleeding
- C Compress with pressure and/or packing
- C Compress with a tourniquet
- √ Wait for help to arrive









## For more information:

STOPTHEBLEED.ORG



The only thing more tragic than a death... is a death that could have been prevented.